

Exact self dual vortices in $d = 3$ Majumdar-Papapetrou dilaton gravity

C.S.Aulakh

Institute of Physics, Bhubaneswar, Orissa 751005, India

Abstract

We show that self-dual Nielsen Olesen (NO) vortices in 3 dimensions give rise to a class of exact solutions when coupled to Einstein Maxwell Dilaton gravity obeying the Majumdar-Papapetrou(MP) relation between gravitational and Maxwell couplings , provided certain Chern-Simons type interactions are present. The metric may be solved for explicitly in terms of the NO vortex function and is asymptotic to Euclidean space with signature $(-1,-1,-1)$. The MP electric field is long range but, strictly speaking, the charge of the vortices is zero since the field dies off as $O(1/r(\ln r)^2)$. The total ADM energy integral of such vortices is *zero*. These peculiarities are due to the nature of the two dimensional Greens function.

In an astonishing recent paper [1] it was shown that the self dual “instantonic” solitons of $4 + 1$ dimensional Yang-Mills (YM) theory and the self dual BPS [2] monopoles of $3 + 1$ dimensional Yang-Mills-Higgs theory give rise to a class of exact stable static solutions even after they are coupled to Einstein-Maxwell(EM) gravity at the Majumdar- Papapetrou(MP) point [3] (i.e the Maxwell coupling is fixed by the gravitational coupling so that the static Newton and Coulomb potentials are equal in magnitude). Important additional requirements are the inclusion of certain non-minimal terms of form $\epsilon^{MNL PQ} A_M F_{NL}^A F_{PQ}^A$ in $4 + 1$ dimensions (and its dimensional reduction in $3 + 1$ dimensions) and coupling to a dilaton in $3 + 1$ dimensions. Here A_M, A_M^A are the Electromagnetic and YM gauge potentials and F_{MN}^A is the YM field strength . These terms are responsible for the soliton acquiring a topological electric charge coupled to the Maxwell field by a mechanism proposed similar to one proposed for $4 + 1$ Yang Mills Chern Simons(YMCS) theory solitons in [4–6]. This electric charge acts as a Bogomol’nyi bound under the ADM [7] mass of field configurations in the theory. The self dual solitonic solutions of Ref.[1] saturate this bound in such a way as to imply the existence of a Killing spinor with respect to a certain Einstein Maxwell covariant derivative. Since the bosonic action used is a subset of the $d = 5, N = 2$ supergravity [8] it follows that the solutions preserve one of the two supersymmetries thus providing yet another example of the supersymmetry associated with self dual solutions of field equations [9]. The result of Ref.[1] allowed us to immediately confirm [10] the conjecture [5,6] concerning the gravitational stabilization of instantonic configurations in $SU(N), N \geq 3$ YMCS theory. The elegance and naturalness of the arguments of Ref.[1] lead one to expect that generalizations to other dimensions and types of solitons (strings, vortices etc.) should exist. In this letter we obtain solutions in 3 dimensions analogous to those of Ref.[1] in four and five dimensions. However the peculiarities of the spherically symmetric Greens function in two spatial dimensions (i.e the logarithm) lead to some peculiar properties for the solutions we generate. The asymptotic spacetime is flat but Euclidean (with no deficit angle). Correspondingly, the total ADM energy integral yields *zero* and the MP electric field, though long range, dies off marginally faster (by a factor of $1/(\log r)^2$) than that of a charged ob-

ject. Thus the expected topological charge is also zero. Nevertheless our solutions appear sufficiently similar to those of Ref.[1] to keep alive the expectation of generalizations to other models where flatspace self-dual solutions exist .

The action we shall treat is the sum of three pieces S_{gr} , S_{SD} and S_{CS} :

$$S_{gr} = -(16\pi G)^{-1} \int d^3x E(R + e^{2b\sigma} F^2 + 2(\partial\sigma)^2) \quad (1)$$

$$S_{SD} = - \int d^3x E \left(\frac{1}{4g^2} e^{2b_a\sigma} f^2 + e^{2b_\psi\sigma} |D\psi|^2 + \frac{g^2}{2} e^{2b_u\sigma} (\psi^\dagger\psi - v^2)^2 \right) \quad (2)$$

$$S_{CS} = \int d^3x \epsilon^{\mu\nu\lambda} A_\mu (\kappa_1 \partial_\nu a_\lambda + \kappa_2 \partial_\nu J_\lambda) \quad (3)$$

Here F, f are field strengths of the MP (A_μ) and NO (a_μ) Abelian gauge potentials, ψ the charged scalar field of the NO model, σ the dilaton field and $J_\lambda = \frac{1}{2i} \psi^\dagger \vec{D}_\lambda \psi$, ($D_\lambda = \partial_\lambda \psi - ia_\lambda \psi$) the NO current. Notice that the MP coupling has been equated to the gravitational one : $g_e^2 = 4\pi G$. The dilaton couplings b, b_a, b_ψ, b_u will be chosen in the course of the calculation to allow an exact solution of the full theory given a flat space self dual NO vortex solution. Note, however, that the different terms in the action do *not* scale in the same way under the rescaling $\sigma \rightarrow \sigma + const$ for any choice of scaling weights for the other fields unless one also scales couplings. The other possible Chern-Simons terms can be added to S_{CS} without affecting our conclusions . Our conventions for gravitational quantities are those of [11] E is the determinant of the *dreibein* while $\epsilon^{\mu\nu\lambda}$ is the 3-d antisymmetric tensor density ($\epsilon^{012} = 1$).

To proceed we note that the MP form of the metric in $D + 1$ dimensions can be written in a form equivalent to that of [12] as ($i = 1, 2$) :

$$ds^2 = -\frac{1}{B^{D-2}} dt^2 + B(x^1, x^2) dx^i dx^i \quad (4)$$

While this form of the metric can also be derived from considerations of the existence of a Killing spinor [1], one can view this ansatz simply as dictated by the need to ensure that the off diagonal terms of the Einstein tensor are automatically zero given that the spatial

metric is conformal to the Euclidean one. In $2 + 1$ dimensions we therefore take the metric to be $Diag(-1, B, B)$. We also write $B = e^{2\phi}$

The field equations are :

$$\tilde{G}_{\mu\nu} = G_{\mu\nu} + 8\pi GT_{\mu\nu}(A, \sigma) = -8\pi GT_{\mu\nu}^{SD}(a_\mu, \psi) \quad (5)$$

$$\partial_\mu(Ee^{2b\sigma}F^{\mu\nu}) = -4\pi G\epsilon^{\nu\mu\lambda}\partial_\mu(\kappa_1 a_\lambda + \kappa_2 J_\lambda) \quad (6)$$

$$\begin{aligned} \partial_\mu(Eg^{\mu\nu}\partial_\nu\sigma) - (b/2)e^{2b\sigma}EF^2 &= 8\pi GE((b_a/4g^2)e^{2b_a\sigma}f^2 \\ &+ b_\psi e^{2b_\psi\sigma} |D\psi|^2 + (g^2/2)b_u e^{2b_u\sigma}(\psi^\dagger\psi - v^2)^2) \end{aligned} \quad (7)$$

$$\partial_\mu(e^{2b_a\sigma}Ef^{\mu\nu}) = -2g^2Ee^{2b_\psi\sigma}J^\nu - g^2\epsilon^{\nu\mu\lambda}(\partial_\mu A_\lambda(\kappa_1 - \kappa_2\psi^\dagger\psi)) \quad (8)$$

$$D_\mu(e^{2b_\psi\sigma}ED^\mu\psi) = e^{2b_u\sigma}E\frac{\partial U}{\partial\psi^\dagger} + i\kappa_2\epsilon^{\nu\mu\lambda}\partial_\nu A_\mu D_\lambda\psi \quad (9)$$

where $U = (g^2/2)(\psi^\dagger\psi - v^2)^2$ is the potential.

When the metric is flat and $\sigma = 0, \kappa_i = 0$ the model reduces to the selfdual Abelian Higgs model whose field equations are solved and the energy minimized provided the fields are static, $a_0 = 0$, and the Bogomol'nyi equations:

$$(D_i \mp i\epsilon_{ij}D_j)\psi = 0 \quad (10)$$

$$f_{ij} = \pm\epsilon_{ij}g^2(\psi^\dagger\psi - v^2) \quad (11)$$

are satisfied. These first order equations may be decoupled to yield the well known vortex equation [13] :

$$\partial^2 \ln \frac{\psi^\dagger\psi}{v^2} = g^2(\psi^\dagger\psi - v^2) + 4\pi \sum_k |n_k| \delta^{(2)}(\vec{r} - \vec{r}_k) \quad (12)$$

where $\{n_k, \vec{r}_k\}$ are the winding numbers and locations (positions of the zeros of ψ) of an ensemble of (anti)self dual vortices.

We now show that with a certain choice of dilaton weights every solution of the above self duality equations gives rise to an *exact, explicit* solution of the field equations of the full theory. We impose the relation :

$$e^{2b_\psi\sigma} = e^{2b_a\sigma} B^{-1} = B e^{2b_u\sigma} \quad (13)$$

which will be satisfied for a certain choice of weights provided $\sigma = \delta\phi$. Then it is easy to see, using the assumed flatspace self duality of the NO fields, that provided

$$\begin{aligned} A_0 &= \mp \frac{E e^{2b_\psi\sigma}}{\kappa_2 B} \\ A_i &= 0 \end{aligned} \quad (14)$$

and $\kappa_1/\kappa_2 = v^2$ the field equations for the fields a_μ, ψ are satisfied. Furthermore it is easy to check that all spatial components of the Einstein tensor and the stress tensor of the matter sector vanish. Hence the spatial components of the Einstein equations are satisfied provided the spatial stress tensor of the fields A_μ, σ vanishes which requires that

$$e^{2b_\psi\sigma} = \frac{s\bar{s}\kappa_2 e^{-b\sigma}}{b} \quad (15)$$

where $s = \pm$ is an independent sign and the reason for being careful about putting $E/B = \text{sign}(B) = \bar{s}$ (although we took $B = e^{2\phi}$!) will become clear *a posteriori*. Thus one immediately has $b_\psi = -b/2, \kappa_2 = \bar{s}sb$.

Since we are looking for a smooth soliton solution with a dispersed energy density we do not expect any zeros of the metric to occur. We therefore assume that the sign of B is constant. This will be justified *a posteriori* and can, in any case, be relaxed (if one wishes to study superpositions of vortices and black holes etc.) by taking the equation we derive to hold piece wise in every region labelled by a given sign of B (since our equation is independent of the sign of B). One finds that three remaining nontrivial field equations (i.e those for G_{00}, A_0, σ) reduce to the *single* flat space equation :

$$\begin{aligned} \partial^2 e^{2\phi} &= \pm(16\pi G)\epsilon_{ij}\left(\frac{v^2 f_{ij}}{2} + \partial_i J_j\right) \\ &\equiv (16\pi G)v^2(g^2(\psi^\dagger\psi - v^2) - \frac{1}{2}\partial^2(\frac{\psi^\dagger\psi}{v^2})) \end{aligned} \quad (16)$$

if and only if one identifies $\sigma = \phi, b = 2$ so that $b_\psi = -1, b_a = 0, b_U = -2$. Now, using the vortex equation, it immediately follows that the regular solution of the above Poisson equation is:

$$e^{2\phi} = -C + \mu \left(\log\left(\frac{\psi^\dagger \psi}{v^2}\right) - \frac{\psi^\dagger \psi}{2v^2} - 2 \sum_k |n_k| \log(v^2 |\vec{r} - \vec{r}_k|) \right) \quad (17)$$

where C is an arbitrary constant and $\mu = 16\pi G v^2 > 0$. Thus it follows that as $r \rightarrow \infty, B \rightarrow -2\mu |N| \log(r); (\sum_k n_k = N)!!$ With suitable C , B is negative everywhere since $\psi^\dagger \psi$ grows from 0 at the locations of the vortices to v^2 at ∞ . A sensible choice of asymptotic coordinates is thus $\tilde{r} = r \sqrt{2\mu |N| \log r}$ which, to leading order in $1/\log r$ converts the metric to a flat metric with signature $(-1, -1, -1)!$. Note that apart from implying that the dilaton field has a constant imaginary part: $Im(\sigma) = i\pi/2$ (which is consistent with the reality of the action etc), there is no other *outré* consequence of our solution.

The ADM “energy” (since our metric has signature $(-1, -1, -1)$ there is really no energy in the usual sense) integral gives

$$\mathcal{E} = - \int d^2 x T_0^0 |B| \equiv - \frac{\bar{s}}{16\pi G} \int d^2 x \partial^2 \phi \quad (18)$$

and thus vanishes given the smoothness and regularity of our solution.

Finally, the asymptotic electric field in the new coordinates is

$$F^{\tilde{r}0} = \mp \frac{s}{4\mu |N| \tilde{r} (\log \tilde{r})^2} \quad (19)$$

so that its surface integral dies logarithmically.

The pathologies of 3 dimensions and the special couplings of our ‘dilaton’ are responsible for the peculiarities of our solution. Nevertheless as an addition to the library of exact solutions of gravitationally coupled theories it merits further investigation and motivates the search for further examples of the mechanism of Ref.[1] which involves as an essential part the acquisition of *topological electric fields* by magnetic configurations in the presence of CS type couplings [4–6,1]. We have also found flatpace (Minkowski) versions of these results (and of those for $d=4+1$ in Ref.[10]) by choosing suitable dilaton couplings.

ACKNOWLEDGMENTS

I am grateful to Avinash Khare for useful and stimulating discussions.

REFERENCES

- [1] G.W.Gibbons, D.Kastor, L.A.J.London, P.K.Townsend and J.Traschen, hep-th/9310118.
- [2] E.B.Bogomol'nyi, Sov. J. Nucl. Phys. **24**,449(1976); M.K. Prasad and C.M. Sommerfield, Phys. Rev. Lett. **35**,760(1975).
- [3] S.D.Majumdar, Phys.Rev.**72**,390(1947); A.Papapetrou, Proc. R. Irish Acad. **A51**,191(1947).
- [4] C.S.Aulakh, Mod. Phys. Lett **A7**,2119(1992).
- [5] C.S.Aulakh, Mod. Phys. Lett. **A7**,2469(1992).
- [6] C.S.Aulakh and V.Soni, Int. J.Mod.Phys. **A8**,1653(1993).
- [7] R.Arnowitt, S.Deser and C.Misner, Phys. Rev. **117**,1595(1960); Phys. Rev. **118**,1100(1960); Phys. Rev. **122**,997(1961).
- [8] M.Günyadin, G.Sierra, and P.K.Townsend, Nucl. Phys. **B242**, 244(1985); *ibid* **B253**, 573(1985).
- [9] E. Witten and D. Olive, Phys. Lett. **B78**,97 (1978); Z.Hlousek and D.Spector, Nucl. Phys. **B370**,143 (1992); Phys. Lett. **B283**,75(1992).
- [10] *Topological electropoles in 4 + 1 dimensional EYMCS theory* , C.S.Aulakh and V.Soni, IP/BBSR/94-11, hep-th 9403085.
- [11] S.Weinberg, *Gravitation and Cosmology*, (Wiley, New York, 1972).
- [12] R.C.Myers, Phys. Rev. **D35**,455(1987).
- [13] A.Jaffe and C.Taubes *Vortices and Monopoles* (Birkhauser, Boston ,1980).